

BLUE N TANK-MIX COMPATIBILITY in CEREALS

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW

Always add BlueN to the tank mix last. Use a minimum water volume of 200 l/ha when using BlueN in tank mix.

This version supersedes all previous editions

BOLD CAPITALS = Physically & Biologically Compatible	<i>Italics</i> = Physically Compatible with Continuous Agitation	Red on Orange = Do not mix as bacterial survival is severely impacted
Normal = Physically Compatible	BOLD ITALICS = Biologically Compatible with Continuous Agitation	RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE

ADJUVANTS				
Activator 90		<i>Codacide</i>		Silwett L-77
Adigor		Dash HC		Vivolt
Biopower		Kantor		

TRACE ELEMENTS				
Epsotop		Kinsidro Grow		Manganese Sulphate

HERBICIDES				
Agritox		Hurricane SC		Pelican Delta
Atlantis OD		Incelo + Biopower		Pixxaro EC
Attribut		Inka SX		Proclus
Axial Pro		Jubilee SX		Proverb + Biopower
Boxer		Leystar		Quelex
Broadway Star		Liberator		Ratio SX
Broadway Ultra + Phase II		Lontrel 600		Shield Pro
<i>Chekker OD</i>		Manhattan + Phase II		Slalom
Counter SX		Memphis + Phase II		Spitfire
Dakota		Monolith		Starane Hi-Load HL
Dingo		Most Micro		Starane XL
Easel		Nevada		Thor
Foxtrot		Omnera LQM		Topik
Galaxy		Pacifica Plus + Biopower		Vixen
Herboxone		Palio		Zypar
Hunter				

FUNGICIDES				
Amistar		Elatus Era + Univoq		Priaxor EC + Revystar XE
Arizona		Fandango		Priaxor EC
Ascra Xpro		Flexity		Proline 275
Aviator Xpro		Folicur		Prosaro
Caramba		Folicur + Talius		Questar
Comet 200		Fontelis		Revystar XE
Custodia		Helix		Rylox
Cyflamid		Imtrex		Takumi SC
Deacon		Kestrel		Talius
Deacon + Talius		Kumulus		Thiopron
Decoy 250EC		Mirage		Univoq
Elatus Era		Monkey		Unix
Elatus Era + Questar		Penncozeb 80WP		Verben

INSECTICIDES				
Clayton Sparta EC		Hallmark Zeon		Sumi-alpha
CythrIn 500 EC		Insyst		Teppeki
Decis Forte		Mavrik		

PLANT GROWTH REGULATORS				
3C Chlormequat 750 (max rate 2.0 l/ha)		Medax Max		Sonis
Canopy		Medax Top		Stabilan 750 (max rate 2.0 l/ha)
Chlormequat + Medax Top		Moddus		Terpal
Chlormequat + Sonis				

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower’s risk. For mixes that are supported physically , no information is available on the efficacy of the tank-mix partner.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

Mixing Notes:

Half fill the tank with clean water start agitation and add the required quantity of Blue N to the spray tank, wait until fully dissolved and dispersed before adding the remaining water. Maintain agitation during topping up the tank and whilst spraying. Spray within 6 hours of mixing Blue N. If using Blue N in tank mixture, then add Blue N to the spray tank last, when the tank is full of water, to allow for maximum dilution.

Changes:

New list; 27 Feb 24 - addition of Palio; 14th November 2024 - addition of Broadway Ultra, Incelo, Manhattan, Proverb and Pacifica Plus with adjuvant; 10th March 25 - addition of crop nutrition products; 14th April 2025 - alteration of Mixing Notes; 22nd April 2025 - update on mixing key

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL

Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK

All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

BlueN contains *Methylobacterium symbioticum* SB23

Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk

BLUE N TANK-MIX COMPATIBILITY FOR USE IN OILSEED RAPE

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW

Always add BlueN to the tank mix last. Use a minimum water volume of 200 l/ha when using BlueN in tank mix.

This version supersedes all previous editions

BOLD CAPITALS = Physically & Biologically Compatible	<i>Italics</i> = Physically Compatible with Continuous Agitation	Red on Orange = Do not mix as bacterial survival is severely impacted
Normal = Physically Compatible	BOLD ITALICS = Biologically Compatible with Continuous Agitation	RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE

ADJUVANTS

Activator 90		Codacide		Vivolt	
Adigor		Dash HC			
Biopower		Silwett L-77			

HERBICIDES

Astrokerb		Fusilade Forte		Panarex	
Belkar		Fusilade Max		Ralos	
Centium CS		Kerb Flo 500		Select Prime	
Centurion Max		Kerb Flo		Shield Pro	
Cleancrop Forward		Korvetto		Sultan 500EC	
Dymid		Laser		Targa Super	
Falcon		Leopard 5 EC		VextaDim 240EC	
Fox		Lontrel 600			

FUNGICIDES

Amistar		Decoy 250EC		Propulse	
Aviator Xpro		Folicur		Proline 275	
Bogard		Kestrel		Prosaro	
Caramba		Mirage		Shepherd	
Custodia		Monkey		Toprex	
Deacon		Plover			

INSECTICIDES

Clayton Sparta EC		Insyst		Sumi-alpha	
Cythrin 500 EC		Karate 2.5 WG		Teppeki	
Decis Forte		Mavrik		Tracer	
Hallmark Zeon		Steward			

PLANT GROWTH REGULATORS

Caryx					
-------	--	--	--	--	--

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower’s risk. For mixes that are supported physically , no information is available on the efficacy of the tank-mix partner.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

Mixing Notes:
Half fill the tank with clean water start agitation and add the required quantity of Blue N to the spray tank, wait until fully dissolved and dispersed before adding the remaining water. Maintain agitation during topping up the tank and whilst spraying. Spray within 6 hours of mixing Blue N. If using Blue N in tank mixture, then add Blue N to the spray tank last, when the tank is full of water, to allow for maximum dilution.

Changes:
New List. 2nd November 2023 - updated with new herbicides and fungicides.; 14th November 2024 - added Astrokerb, Kerb Flo 500, Plover and Shepherd; 14th April 2025 - adjustment of Mixing Notes; 22nd April 25 - updated mixing key

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL
Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK
All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
BlueN contains *Methylobacterium symbioticum* SB23
Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk

BLUE N TANK-MIX COMPATIBILITY FOR USE IN POTATOES

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW

Always add BlueN to the tank mix last. Use a minimum water volume of 200 l/ha when using BlueN in tank mix.
This version supersedes all previous editions

BOLD CAPITALS = Physically & Biologically Compatible	<i>Italics</i> = Physically Compatible with Continuous Agitation	Red on Orange = Do not mix as bacterial survival is severely impacted
Normal = Physically Compatible	BOLD ITALICS = Biologically Compatible with Continuous Agitation	RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE

ADJUVANTS				
Activator 90		Biopower		Dash HC
Adigor		Codacide		Silwett L-77

HERBICIDES				
Basagran SG		Leopard 5 EC		Shotput
Falcon		Most Micro		Titus + Codacide
Fusilade Max		Panarex		Titus + Phase II
Laser		Sencorex Flo		Titus + Vivolt

FUNGICIDES				
Amistar		Nautile DG + Zorvec Endavia		Ranman Top
Caligula		Nautile DG + Zorvec Entecta		Revus
Carial Flex		Option		Signum
Enervin		Option + Proxanil + Zorvec Endavia		Vendetta
Fubol Gold		Option + Zorvec Endavia		Versilus
Gachinko		Option + Proxanil + Zorvec Entecta		Zorvec Endavia
Infinito		Option + Zorvec Entecta		Zorvec Enicade
Lieto		Penncozeb 80WP		Zorvec Entecta
Manzate 75WG + Zorvec Endavia		Percos		Zorvec Entecta + Phase II
Manzate 75WG + Zorvec Entecta		Proxanil		

INSECTICIDES				
Clayton Sparta EC		Decis Forte		Sumi-alpha
CythrIn 500 EC		Insyst		Teppeki (Seed crops only)

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower’s risk. For mixes that are supported physically , no information is available on the efficacy of the tank-mix partner.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

Mixing Notes:
Half fill the tank with clean water start agitation and add the required quantity of Blue N to the spray tank, wait until fully dissolved and dispersed before adding the remaining water. Maintain agitation during topping up the tank and whilst spraying. Spray within 6 hours of mixing Blue N. If using Blue N in tank mixture, then add Blue N to the spray tank last, when the tank is full of water, to allow for maximum dilution.

Changes:
New List; 2nd November 2023 - additions to herbicides and fungicides; 14 th November 2024 - addition of new herbicides and fungicides, including Zorvec Entecta; 10th March 25 - addition of Enervin; 14th April 2025 - adjustment of Mixing Notes; 22nd April

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL
Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK
All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
BlueN contains *Methylobacterium symbioticum* SB23
Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk

BLUEN TANK-MIX COMPATIBILITY FOR USE IN SUGAR BEET

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW

Always add BlueN to the tank mix last. Use a minimum water volume of 200 l/ha when using BlueN in tank mix.
This version supersedes all previous editions

BOLD CAPITALS = Physically & Biologically Compatible	<i>Italics</i> = Physically Compatible with Continuous Agitation	Red on Orange = Do not mix as bacterial survival is severely impacted
Normal = Physically Compatible	BOLD ITALICS = Biologically Compatible with Continuous Agitation	RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE

Adjuvants				
Activator 90		Biopower		Dash HC
Adigor		<i>Codacide</i>		Silwett L-77

Herbicides				
Betasana SC		Fusilade Max		Panarex
Bettix WG		Goltix 70SC		Shield Pro
Debut		Laser		Targa Super
Falcon		Leopard 5 EC		Venzar 500SC
Fusilade Forte		Lontrel 600		

Fungicides				
Revystar XE		Thiopron		

Insecticides				
Clayton Sparta EC		Decis Forte		Insyst
CythrIn 500 EC		Hallmark with Zeon		Teppeki

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower’s risk. For mixes that are supported physically , no information is available on the efficacy of the tank-mix partner.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Technical Hotline on 0800 689 8899

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, PH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Spray able with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

Mixing Notes:

Half fill the tank with clean water start agitation and add the required quantity of Blue N to the spray tank, wait until fully dissolved and dispersed before adding the remaining water. Maintain agitation during topping up the tank and whilst spraying. Spray within 6 hours of mixing Blue N. If using Blue N in tank mixture, then add Blue N to the spray tank last, when the tank is full of water, to allow for maximum dilution.

Changes:
2nd November - additional herbicides, fungicides and insecticides added.14th November 2024 - addition of Debut and Venzar 500SC; 14th April 25 -adjustment of Mixing Notes; 22nd April 2025 - updated mixing key

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL
Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK.
All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
BlueN contains *Methylobacterium symbioticum* SB23
Technical Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk

BLUE N TANK-MIX COMPATIBILITY FOR USE IN MAIZE

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW

Always add BlueN to the tank mix last. Use a minimum water volume of 200 l/ha when using BlueN in tank mix.
This version supersedes all previous editions

BOLD CAPITALS = Physically & Biologically Compatible	<i>Italics</i> = Physically Compatible with Continuous Agitation	Red on Orange = Do not mix as bacterial survival is severely impacted
Normal = Physically Compatible	BOLD ITALICS = Biologically Compatible with Continuous Agitation	RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE

ADJUVANTS					
Activator 90		Codacide		Vivolt	
Adigor		Dash HC			
Biopower		Silwett L-77			

HERBICIDES					
Accent		Dragster + Vivolt		Oceal	
Botiga		Dual Gold		Peak	
Callisto		Elumis		Principal Forte + Codacide	
Callisto + Dragster		Herboxone		Principal Forte + Vivolt	
Callisto + Maister		Laudis		Rosan	
Callisto + Principal Forte		Leystar		Shield Pro	
Camix		Lontrel 600		Starane Hi-Load HL	
Choriste		Maister		Titus	
Dingo		Most Micro		Wing P	
Dragster + Codacide		Nico Pro 4SC			

INSECTICIDES					
CythrIn 500 EC					

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower’s risk. For mixes that are supported physically , no information is available on the efficacy of the tank-mix partner.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

Mixing Notes:
Half fill the tank with clean water start agitation and add the required quantity of Blue N to the spray tank, wait until fully dissolved and dispersed before adding the remaining water. Maintain agitation during topping up the tank and whilst spraying. Spray within 6 hours of mixing Blue N. If using Blue N in tank mixture, then add Blue N to the spray tank last, when the tank is full of water, to allow for maximum dilution.

Changes:
New List: 2nd Nov23 - updated list with new herbicides and insecticide; 14th November 2024 - addition of Dragster and Principal Forte mixes.10th March 25 - addition of Botiga; 14th April 25 - adjustment of Mixing Notes; 22nd April 25 - updated mixing key; 12th May

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL
Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK.
All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
BlueN contains *Methylobacterium symbioticum* SB23
Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk

BLUE N TANK-MIX COMPATIBILITY FOR USE IN GRASS

PLEASE ENSURE YOU READ THE NOTES SECTION BELOW

Always add BlueN to the tank mix last. Use a minimum water volume of 200 l/ha when using BlueN in tank mix.
This version supersedes all previous editions

BOLD CAPITALS = Physically & Biologically Compatible	<i>Italics</i> = Physically Compatible with Continuous Agitation	Red on Orange = Do not mix as bacterial survival is severely impacted
Normal = Physically Compatible	BOLD ITALICS = Biologically Compatible with Continuous Agitation	RED CAPITAL ON BLUE = CROP INJURY - INCOMPATIBLE

HERBICIDES					
Cabadex		Envy		Shield Pro	
Dingo		Leystar		Sickle	
Doxstar Pro		Kyleo		Thistlex	
Easel		ProClova XL			

This summary sheet provides information on tank-mixes conducted with this product. Corteva Agriscience™ gives no warranty or accepts no liability in respect to compatibilities that are beyond the label. This additional information is for guidance purposes only and use is at grower’s risk. For mixes that are supported physically , no information is available on the efficacy of the tank-mix partner.

Important Note: Do not use any tank-mix if the mixture cannot be applied within the recommendations for individual products or if specific tank-mix restrictions apply.

Larger, more complicated mixes; unlisted mixes or replacing alternative products to those listed in this table are not endorsed. However, for further information or to register a tank-mix request please contact your local Corteva representative or call the Hotline on 0800 689 8899.

Physical compatibility data is based on the results of an industry recognised standard test. Be aware that in the field, water hardness, pH, temperature etc. can vary. Biological compatibility data are based on a small number of field trials, Corteva Agriscience™ cannot warrant the efficacy of other tank-mix partners. Where only Physical Compatibility or Sprayable with Continuous Agitation is shown, Corteva Agriscience™ has no biological data regarding efficacy or crop safety and only has data on physical compatibility, so growers use these mixes at their own risk.

Mixing Notes:
Half fill the tank with clean water start agitation and add the required quantity of Blue N to the spray tank, wait until fully dissolved and dispersed before adding the remaining water. Maintain agitation during topping up the tank and whilst spraying. Spray within 6 hours of mixing Blue N. If using Blue N in tank mixture, then add Blue N to the spray tank last, when the tank is full of water, to allow for maximum dilution.

Changes:
New List; 10th March 25 - addition of Proclova XL; 14th April 25 - adjustment of Mixing Notes; 22nd April updated mixing key

USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.
FOR FURTHER INFORMATION INCLUDING WARNING PHRASES AND SYMBOLS PLEASE REFER TO THE PRODUCT LABEL
Corteva Agriscience™, Unit H4, Building H, Melbourn Science Park, Cambridge Road, Melbourn, Cambridgeshire, SG8 6HB, UK
All other brand names are trademarks of other manufacturers for which proprietary rights may exist.
BlueN contains *Methylobacterium symbioticum* SB23
Hotline: 0800 689 8899 | ukhotline@corteva.com | www.corteva.co.uk